

Construction of extremal local positive operator-valued measures under symmetry

S. Virmani^{1,2} and M.B. Plenio²

¹ Gruppo di Ottica Quantistica, Dip. di Fisica 'A. Volta', Università degli Studi di Pavia, via Bassi 6, I-27100 Pavia, Italy

² QOLS, Department of Physics, Blackett Laboratory, Imperial College, Prince Consort Road, London SW7 2BW, UK
(February 1, 2008)

We study the local implementation of POVMs when we require only the faithful reproduction of the statistics of the measurement outcomes for all initial states. We first demonstrate that any POVM with separable elements can be implemented by a separable super-operator, and develop techniques for calculating the extreme points of POVMs under a certain class of constraint that includes separability and PPT-ness. As examples we consider measurements that are invariant under various symmetry groups (Werner, Isotropic, Bell-diagonal, Local Orthogonal), and demonstrate that in these cases separability of the POVM elements is equivalent to implementability via LOCC. We also calculate the extrema of these classes of measurement under the groups that we consider, and give explicit LOCC protocols for attaining them. These protocols are hence optimal methods for locally discriminating between states of these symmetries. One of many interesting consequences is that the best way to locally discriminate Bell diagonal mixed states is to perform a 2-outcome POVM using local von Neumann projections. This is true regardless of the cost function, the number of states being discriminated, or the prior probabilities. Our results give the first cases of local mixed state discrimination that can be analysed quantitatively in full, and may have application to other problems such as demonstrations of non-locality, experimental entanglement witnesses, and perhaps even entanglement distillation.

PACS numbers: 03.67.-a, 03.67.Hk

I. INTRODUCTION

The nature of local and global quantum operations plays a key role in Quantum Information theory [1]. However, although global quantum operations are well characterised [2], there is still no convenient mathematical classification of the so called LOCC transformations (operations that can be implemented locally with classical communication). In recent years some progress has been made by trying to understand the non-local properties of unitaries and by trying to implement operations with a minimum amount of prior entanglement [3,4,6]. Nevertheless, a more general answer remains elusive, and so alternative classes of operations have been defined that can be more easily characterised. Perhaps the most interesting examples are the Separable and PPT operations introduced by Rains [7,8]. These classes of operations (see section II for definitions) are mathematically more tractable than the LOCC class, and have been particularly useful as tools for investigating the entanglement of distillation and local state discrimination [8,9]. They have also been used to extend the conventional definitions of entanglement measures [10,11], giving indications that quantum state transformations can adopt a simplified structure under the class of PPT operations [12].

The separable and PPT operations will also play an important role in this work, where we investigate the local implementation of quantum measurements. This question is significant as in many protocols for quantum communication it is important to know how a distantly separated Alice and Bob may be able to gain information about quantum states using only local operations and classical communication. Examples include entanglement distillation and cryptographic scenarios [9,10,13]. The issue of local measurements is also important

for efforts to locally infer some form of entanglement, such as in developing tests of non-locality, or the local detection of inseparability [14]. However, there is perhaps a much more fundamental reason for the investigation of local measurements: in all uses of quantum states, all we ever 'see' at the end of the day are measurement outcomes, so one of the most important things to know is what kinds of statistics are possible according to the classes of operations that we are restricted to. As the LOCC operations are an important paradigm in quantum information, it is important to know what kinds of measurement outcomes we can obtain with them.

Most of the discussion of local measurements in recent years has been directed particularly towards local state discrimination [15–19]. However, in this work we pay more attention to the actual implementation of measurements locally. We will be inspired by initial steps in this direction taken by [9,20], where both the concepts of PPT-operations and symmetry were used to design a certain type of cryptographic scheme for Werner states of dimensions $2^n \times 2^n$ (work that was extended to the multiparty setting in [21]). Unlike those investigations, however, we will not be directly interested in any particular application or cost functions, as our main aim will be to obtain techniques for deciding the local implementability of measurements, with particular regard to finding extreme points. Some of the methods are quite general, and have implications for finding extremal POVMs under constraints other than local implementability.

This paper is structured as follows. In section II we discuss the connection between the entanglement of POVM elements and implementability via the different classes of operations introduced by Rains [8]. In sections III and IV we use these ideas to derive the entire (convex) set of local measurements that satisfy Isotropic and Werner symmetries, also giving the

extreme points. These particular symmetries are amenable to an elegant form of solution that is not possible for more complicated symmetry groups, where we will require stronger methods. So in section V we discuss general techniques for deriving the extrema of POVMs under a quite general class of constraint that includes the constraints of separability and PPT-ness. In section VI we apply these techniques to Bell and OO [22] symmetries, deriving the sets of locally implementable measurements and their extreme points. The Bell symmetries are particularly interesting, as we show that the best way to locally discriminate Bell diagonal states is just to perform a 2-outcome POVM using local von Neumann projections. This is true regardless of the cost function, the number of states being discriminated, or the prior probabilities. Finally in section VII we summarise and discuss the implications of our results.

II. ENTANGLEMENT CLASSIFICATION OF POVM ELEMENTS AND RELATIONSHIP TO CLASSES OF OPERATIONS

We begin by investigating the relationship between the entanglement properties of POVM elements and the resources required to implement the POVM. The key result of this section will be the demonstration that the separability (PPT-ness) of the POVM elements is equivalent to the implementability of the corresponding POVM by separable (PPT) operations.

Let us begin by reviewing the concepts of PPT and Separable operations. The set of PPT operations P is the set of completely positive (CP) transformations that remain completely positive when conjugated with the partial transposition operator Γ , i.e. both P and $\Gamma \circ P \circ \Gamma$ are completely positive. It is irrelevant whether the partial transposition is taken as transposition of Bob's or Alice's system, as long as the choice is fixed. The set of Separable operations is the class of operations S that can be written as:

$$S(\rho) = \sum_n A_n \otimes B_n \rho A_n^\dagger \otimes B_n^\dagger. \quad (1)$$

It is not too difficult to verify [7] that all LOCC operations are separable, and all separable operations are PPT. An example of a CP operation that is not PPT is the creation of an NPT state from a PPT one [7]. The PPT and Separable operations can also be characterised in an interesting way in the light of the Jamiolkowsky isomorphism [5–9,20].

This classification of operations has allowed interesting bounds to be derived in the distillation and creation of entangled states [8,23], and is also useful for the study of local quantum state discrimination [9,20]. In the following, we will be interested particularly in the question of when a measurement procedure can be implemented by LOCC. We will not be concerned with the residual states left over after any measurement process, so a POVM will be considered to be 'locally implementable' if we can perform a measurement using only

local operations and classical communication that allows exactly the same statistical inferences to be made. It is important to note the differences between our use of the word 'local' and that of Beckman et. al. [25] and Groisman et. al. [24]. Those authors are motivated by the implications of quantum operations and observations for causality, and hence they do allow the use of prior entanglement but disallow the use of classical communication.

In our context the classification introduced by [7,8] has a particularly elegant structure, as the PPT and separable definitions of operations directly map to the definitions of PPT and separable states [26,27]. A given POVM described by a set of elements $\{M_k | k = 1..N, M_k \geq 0, \sum M_k = \mathbb{1}\}$ can be implemented by a PPT operation iff all the elements (after normalisation) correspond to PPT states, and can be implemented by a separable operation iff all the elements correspond to separable states:

$$\text{can do POVM by Separable ops} \Leftrightarrow \text{elements separable} \quad (2)$$

$$\text{can do POVM by PPT ops} \Leftrightarrow \text{elements PPT} \quad (3)$$

The fact that all measurements implemented by separable operations must involve separable POVM elements was noted in [9]. The fact that a measurement can be implemented by a PPT operation iff the POVM elements are PPT was noted by [20,28]. As far as we are aware, however, the fact that all POVMs with separable elements can be implemented by a separable operation has not been noted anywhere despite its relatively simple proof. For completeness we therefore present the following proof of correspondence (2):

Observation 1. A given POVM can be implemented by a separable superoperator iff the POVM elements associated with all N outcomes $\{M_k | k = 1..N, M_k \geq 0, \sum M_k = \mathbb{1}\}$ correspond to separable states, i.e. the POVM can be implemented by a separable superoperator iff the density matrices defined by $M_k / \text{tr}\{M_k\}$ are separable.

Proof. Any POVM can be realised as a superoperator where the subscripts n on the Kraus operators have been recorded classically. Any separable operation will correspond to POVM elements where the subscripts n will be grouped into sets S_k corresponding to the different inferences k . Hence the expectation values of any POVM element M_k can be written:

$$\text{tr}\{M_k \rho\} = \sum_{n \in S_k} \text{tr}\{A_n \otimes B_n \rho A_n^\dagger \otimes B_n^\dagger\} \quad (4)$$

$$= \sum_{n \in S_k} \text{tr}\{A_n^\dagger A_n \otimes B_n^\dagger B_n \rho\}. \quad (5)$$

Hence we can write that:

$$M_k = \sum_{n \in S_k} A_n^\dagger A_n \otimes B_n^\dagger B_n \quad (6)$$

As this sum is completely in terms of positive operators, we have that $\text{tr}\{M_k\} = \sum_n \text{tr}\{A_n^\dagger A_n\} \text{tr}\{B_n^\dagger B_n\}$. This means that we can also write:

$$\begin{aligned}\rho_k &:= \frac{M_k}{\text{tr}\{M_k\}} \\ &= \sum_{n \in S_k} \frac{\text{tr}\{A_n^\dagger A_n\} \text{tr}\{B_n^\dagger B_n\}}{\text{tr}\{M_k\}} \frac{A_n^\dagger A_n}{\text{tr}\{A_n^\dagger A_n\}} \otimes \frac{B_n^\dagger B_n}{\text{tr}\{B_n^\dagger B_n\}}.\end{aligned}$$

Hence we see that each POVM element from a separable operation must correspond to a separable state. To see the converse, simply work backwards through the above procedure, using the fact that any positive operator X has a decomposition $X = Q^\dagger Q$. The generalisation to multi-party systems is straightforward $\square$.

Observation 1 is extremely powerful, and can be used to provide simple derivations of many results. For example we can immediately see that an entangled pure state cannot be perfectly locally discriminated from its orthogonal complement, as this would require a POVM with non-separable elements (see also [18]). A similar argument also shows that we cannot perfectly locally distinguish an Unextendible Product Basis (UPB) [29] from an equal mixture of the pure states in its orthogonal complement, adding to the other intriguing discrimination properties of such bases [29].

Is it possible that all separable discrimination protocols can be implemented locally? Unfortunately this is definitely not the case, as has been observed from the example of non-locality without entanglement presented in [30] (see also [16]). The authors of that paper present sets of orthogonal separable states that cannot be discriminated perfectly using only local operations. If all separable POVMs could be implemented locally then those sets of orthogonal states could be discriminated perfectly using only local operations. As this is not the case, this means that not all separable operations can be implemented locally.

Nevertheless, the constraint that any local POVMs must be separable is still quite a strong one, and in some simple cases of high symmetry we will see that it is exactly equivalent to the local implementability of POVMs. The results thereby allow the construction of optimal local discrimination protocols for any number of states of such symmetries under any cost function for any prior probabilities. The ideas may also have other applications in cryptographic schemes, demonstrations of non-locality, and the local detection of entanglement. We will begin by considering the *Isotropic* symmetries.

III. LOCAL OBSERVATIONS WITH ISOTROPIC SYMMETRIES

In this section we begin by analysing the so called *Isotropic* symmetries. The class of measurements invariant under this group is actually a subset of the *OO* symmetric measurements that we will consider later in the paper. However, we consider the *Isotropic* and *Werner* cases independently in this section and the next, as they afford a particularly elegant form of solution and demonstrate the usefulness of the partial transposition mapping between *Isotropic* and *Werner* states.

The *Isotropic* states are those that commute with all local unitaries of the form $U \otimes U^*$, where the $*$ denotes complex conjugation in a fixed local basis. Any isotropic state $\sigma(f)$ on $C^d \otimes C^d$ can be written as:

$$\sigma(f) = f|+\rangle\langle+| + (1-f)\frac{\mathbb{1} - |+\rangle\langle+|}{d^2 - 1}, \quad (7)$$

where $f \in [0, 1]$, and $|+\rangle$ is a canonical maximally entangled state $|+\rangle = \sum |ii\rangle/\sqrt{d}$. In the local observation of such states we can use standard symmetry arguments to restrict our attention to POVMs with elements of the form:

$$M_k = a_k|+\rangle\langle+| + b_k(\mathbb{1} - |+\rangle\langle+|). \quad (8)$$

As we require these elements to form a valid POVM, they must satisfy the constraints:

$$\begin{aligned}a_k &\geq 0 \\ b_k &\geq 0 \\ \sum_k a_k &= 1 \\ \sum_k b_k &= 1.\end{aligned} \quad (9)$$

It is well known that isotropic states are separable iff they are PPT (see e.g. [22]), and so it is easy to compute the additional constraint that each M_k must be separable from $M_k^\Gamma \geq 0$:

$$(d+1)b_k \geq a_k. \quad (10)$$

We will now see that there is a local protocol that can be used to attain any *Isotropic* POVM satisfying equations (9) and (10). The protocol consists of two steps:

- Alice and Bob perform an isotropic twirl [31], $T(\rho)$. This step is superfluous if the states to be observed are isotropic anyway. It is only included to make the total measurement exactly equal to the POVM satisfying (9) and (10).
- Then Alice and Bob perform a measurement with elements N_k described by:

$$N_k = \sum_{i=1}^d |i\rangle\langle i| \otimes [x_k|i\rangle\langle i| + y_k(\mathbb{1}_B - |i\rangle\langle i|)] \quad (11)$$

with

$$\begin{aligned}x_k &= a_k \\ y_k &= \frac{(d+1)b_k - a_k}{d},\end{aligned} \quad (12)$$

where $|i\rangle$ is a pure state from the computational bases of Alice & Bob, and $\mathbb{1}_B$ refers to the identity on Bob's space.

The probability that Alice and Bob will find outcome k from the above procedure given an input state ρ is given by:

$$\text{tr}\{N_k T(\rho)\} = \text{tr}\{T(N_k)\rho\}. \quad (13)$$

It is not difficult to verify that $M_k = T(N_k)$ for all k , and so our two step protocol gives exactly the same statistics for each k as the original measurement. Moreover, examination of the form of the N_k shows that they correspond to a local measurement involving only one way communication, as long as the condition (10) holds (otherwise the N_k will not be positive). Therefore any POVM consisting of PPT Isotropic elements can be attained locally. The transformation above also gives a direct derivation of the extreme points of this class of POVMs. The mapping from the PPT (a_k, b_k) to the (x_k, y_k) is linear and invertible. The x_k and y_k form two independent probability distributions, and so the extreme points will just correspond to situations when the x_k, y_k take on values of $\{0, 1\}$. This observation allows relatively straightforward optimisation of local discrimination of isotropic states under essentially all cost functions.

It is also worthwhile noting that there are many other possible local protocols that can match the PPT isotropic POVMs. This is because the above protocol essentially makes full use of the maximally entangled component of the isotropic states to leave Bob in a residual mixture of some pure state with the identity, which can then be used to discriminate between the original isotropic states. Indeed, any measurement that Alice can perform on a maximally entangled state that would leave Bob with a known pure state can be used to construct a similar protocol. One example is teleportation. Suppose Alice and Bob first teleport a known pure state $|\psi\rangle$ using whichever of the isotropic states that they share. Then if Bob performs a POVM defined by elements $x_k|\psi\rangle\langle\psi| + y_k(\mathbb{1}_B - |\psi\rangle\langle\psi|)$ then they will also achieve the same POVMs as the N_k defined above.

IV. LOCAL OBSERVATIONS WITH WERNER SYMMETRIES

Having calculated the set of locally implementable isotropic POVMs, let us see why we can almost immediately write down the set of locally implementable Werner POVMs [26]. The *Werner* states are those that commute with all local unitaries of the form $U \otimes U$. Any Werner symmetric POVM can be written with elements of the form:

$$M_k = a_k P_A + b_k P_S, \quad (14)$$

where P_A and P_S are the so called *antisymmetric* and *symmetric* projectors respectively [26].

The argument here follows from the partial transposition mapping between Isotropic and Werner operators that was discussed in [22]. The authors of that paper point out that if a certain class of operators is invariant under a given group of local unitary operations $U(\gamma) \otimes V(\gamma)$, then the partial transpositions of those operators will be invariant under the ‘partially conjugated’ group $U(\gamma) \otimes V^*(\gamma)$ (modulo a problem of phases, see [22]). This means that if we have a set of PPT

POVM elements that is invariant under one local symmetry group,

$$\{M_k | \sum M_k = \mathbb{1}, M_k \geq 0, M_k^\Gamma \geq 0\} \quad (15)$$

then the partial transposition of these elements will form another POVM that is a PPT POVM invariant under the partially conjugated group, and vice versa. Indeed the extremal PPT measurements under one group of local unitaries are in one-to-one correspondence with extremal PPT measurements under the ‘partially conjugated’ group:

Theorem 2. If $\{M_k\}$ forms an extremal PPT POVM for one local symmetry group, then the partial transposition $\{M_k^\Gamma\}$ forms an extremal PPT POVM for the partially conjugated symmetry group.

Proof. If a POVM $\{M_k | k = 1 \dots n\}$ is PPT and *not* extremal, then it is possible to write:

$$\{M_k | k = 1 \dots n\} = \{pM_k^1 + (1-p)M_k^2 | k = 1 \dots n\} \quad (16)$$

for some probability p , where $\{M_k^1 | k = 1 \dots n\}$ and $\{M_k^2 | k = 1 \dots n\}$ are also PPT POVMs. Therefore the partial transposition of this equation is also true:

$$\{(M_k)^\Gamma | k = 1 \dots n\} = \{p(M_k^1)^\Gamma + (1-p)(M_k^2)^\Gamma | k = 1 \dots n\}, \quad (17)$$

where $\{(M_k^1)^\Gamma | k = 1 \dots n\}$ and $\{(M_k^2)^\Gamma | k = 1 \dots n\}$ are also PPT POVMs. So the partial transposition of a PPT POVM is extremal iff the PPT POVM itself is extremal $\square$.

This convenient partial transposition connection is precisely the relationship between PPT POVMs of Werner symmetry and PPT POVMs of Isotropic symmetries. Hence all implications discussed above for the local discrimination of isotropic states also immediately apply to local discrimination of Werner states. In particular, the partial transposition of the local protocols given for Isotropic symmetries above will allow us to obtain any PPT POVM with Werner symmetries. To be explicit, let us denote a Werner-symmetric POVM by elements:

$$M_k = a_k P_A + b_k P_S. \quad (18)$$

The condition that the POVM be PPT forces us to impose the requirement that:

$$b_k \left(\frac{d+1}{d-1} \right) \geq a_k \quad (19)$$

in addition to the other positivity and completeness constraints. Any such POVM can be attained by the partial transposition of a PPT isotropic measurement, and so can be attained by the ‘partial transposition’ of the local measurement used to attain that particular isotropic POVM, i.e.:

- First Alice and Bob perform a Werner twirl [31], $T(\rho)$. Again, this step is superfluous if the states to be observed are Werner anyway.

- Then Alice and Bob perform a measurement with elements N_k described by:

$$N_k = \sum_{i=1}^d |i\rangle\langle i| \otimes [x_k|i\rangle\langle i| + y_k(\mathbb{1}_B - |i\rangle\langle i|)] \quad (20)$$

with

$$\begin{aligned} x_k &= \frac{(1-d)a_k + (d+1)b_k}{2} \\ y_k &= a_k. \end{aligned} \quad (21)$$

It can readily be verified that the protocol matches any PPT Werner POVM.

In the other symmetry groups that we will consider in this paper, no elegant and simple solution along the lines of the one found for the Isotropic/ Werner states can work. The reasons for this will be discussed in more detail in the section on OO symmetries and in the appendix. Consequently, for other symmetry groups we have to adopt a different approach. Here we choose to construct the extremal points of the (convex) set of PPT POVMs for the groups that we consider, and then try to find local protocols that match these extrema. We then need to invoke a theorem of convex analysis which states that any convex compact set is the (closed) convex hull of its extreme points [32], and hence if we can show that the extremal points of the PPT measurements can be obtained locally, then by convexity so can the whole set (for a finite number of outcomes the set of PPT POVMs is clearly compact). In the cases that we consider here our search for local protocols will always be successful. In fact, the most difficult exercise is usually the construction of the extremal points. The task is made tractable by some general techniques that we will discuss next.

V. GENERAL TECHNIQUES AND NOTATION

In this section we will explain the major tools that we will use throughout the rest of the paper. Sometimes we will describe these techniques in quite general language, as there may be applications to other problems.

The first technique, that we will also use later in this subsection, is a quite common method for determining constraints on the extrema of convex sets. Suppose that we have a convex set A , and we are trying to decide whether a candidate element s of the set is extremal. When we write the phrase ‘by the following perturbation’, this will mean that we consider the following common approach to deciding extremality. We can consider perturbations of s by a small amount δ . We then try to decompose the candidate extremum s in the following way:

$$\begin{aligned} s &= \frac{1}{2}s(+) + \frac{1}{2}s(-) \\ s(+) &= s + \delta \\ s(-) &= s - \delta. \end{aligned} \quad (22)$$

In order that s be extremal, we will require that at least one of $s(+)$ and $s(-)$ be outside the set A . In our specific application, we will usually choose δ such that the separability and completeness of the POVM is maintained, and so we will have to infer that either $s(+)$ or $s(-)$ be non-positive, and this will allow us to derive strong constraints of the form of the extrema. So when we write ‘by the perturbation δ we can see that the extremal points must be of the form...’ we are in fact referring to the inferences that can be made by the above process.

Now let us consider how we can use some of the structure of sets of separable POVMs in order to simplify the hunt for extremal points. The most important feature that we will rely on is the fact that the properties of positivity, separability and PPT-ness are closed under linear combinations with non-negative scalars. Some of the methods that we will employ can be used to extremise POVMs for any other constraints that have this property, and not only PPT-ness or separability. In light of this we will find it convenient to make the following general definition:

Definition 3. A constraint on two operators M_1, M_2 will be called *homogeneous* if whenever M_1 and M_2 satisfy the constraint then so does $\alpha M_1 + \beta M_2$, for all non-negative scalars α, β . Hence both Separability and PPT-ness are homogeneous constraints. We will sometimes use the symbol $\mathbf{H}$ to denote *any* set of homogeneous constraints under consideration, not necessarily only PPT-ness or separability.

Definition 4. A POVM $\{M_k | k = 1 \dots n, M_k \geq 0, \sum M_k = \mathbb{1}\}$ satisfying any extra *homogeneous* constraints $\mathbf{H}$ under consideration will be called a *feasible* measurement.

The set of feasible POVMs is clearly a convex set itself, in that if $\{M_k^1 | k = 1 \dots n\}$ and $\{M_k^2 | k = 1 \dots n\}$ are feasible POVMs, then so is $\{pM_k^1 + (1-p)M_k^2 | k = 1 \dots n\}$ where $p \in [0, 1]$. Hence we can ask questions about what the extremal POVMs are, and how to find them. One particularly important result in this direction is the following:

Theorem 5. In any extremal feasible POVM, the non-zero operator elements must be linearly independent.

Proof. The proof can be adapted straightforwardly from a proof of this statement for global POVMs by Lo Presti & D’Ariano [33], and indeed their proof is essentially true for *any* constraints that are closed under multiplication by non-negative scalars $\square$.

This theorem will be particularly useful as it will allow us to restrict the number of non-zero outcomes that we need to consider. It also provides another reason why in the case of Isotropic/Werner symmetries the extremal PPT POVMs have essentially two non-zero outcomes, simply because under these symmetries there are at most two non-zero linearly independent POVM elements.

The consideration of homogeneous constraints [34] will allow us to use a technique that involves what we call ‘basic vectors’. This should not be confused with the term ‘basic solution’ that is often used in linear programming problems. In

the rest of this section we will explain exactly what we mean by a ‘basic vector’. Let us suppose that we can find a class of elements $\{V(\alpha)\}$ where α represents one or more parameters (discrete or continuous) with the following properties:

- Each element from $\{V(\alpha)\}$ is non-negative.
- Each element from $\{V(\alpha)\}$ satisfies the homogeneous constraints $\mathbf{H}$ under consideration.
- All POVM elements M_k satisfy the homogeneous constraints under consideration iff they can be written in the following way:

$$M_k = \sum_i p_i^k V(\alpha_i) \quad (23)$$

where the p_i^k are non-negative real numbers.

Then the set $\{V(\alpha)\}$ will be referred to as a set of *basic vectors*. The coefficient p_i^k will be referred to as the *weight* of the basic vector $V(\alpha_i)$ in M_k . Of course this decomposition, and hence the weights, need not be unique. Such sets of basic vectors can be very useful tools for deriving the set of extreme points, and in subsequent sections we will prove a few lemmas that will demonstrate their power. The source of this power is the fact that in the ‘basic vector’ approach, each feasible POVM element is automatically decomposed into a sum (with non-negative weights) of operators that automatically satisfy the homogeneous constraints that we are considering. So we are free to try to perturb any non-zero weights by small amounts without violation of our constraints. This allows us to derive strong limitations on which weights can be non-zero in an extremal measurement. In particular the following two Lemmas prove to be useful:

Lemma 6. In an extremal feasible POVM, no basic vector can have non-zero weight in more than one element M_k of the POVM.

Proof. This can be proved along similar lines to the proof of the following lemma $\square$.

Lemma 7. If a feasible POVM is extremal, then any set of linearly dependent basic vectors must have non-zero weights in at most one element M_k .

Proof. Suppose that we have a set of basic vectors $\{V_m | m = 1 \dots L\}$ such that:

$$\sum_m \lambda_m V_m = 0 \quad (24)$$

for some set of non-zero λ_m ’s. We can take the Hermitian conjugate of this equation and average it with the original to give:

$$\sum_m \frac{1}{2} \lambda_m V_m + \frac{1}{2} \lambda_m^* V_m = \sum_m \text{Re}\{\lambda_m\} V_m = 0. \quad (25)$$

Hence the λ_m s can be taken to be real without loss of generality (this is unless all the original λ_m are purely imaginary,

but in that case we can just divide the equation through by $\sqrt{-1}$). Suppose that the basic vector V_L has non-zero weight in element M_2 , whereas the other basic vectors of the set have non-zero weight in M_1 . Then we have that:

$$\sum_{m \neq L} \frac{\lambda_m}{\lambda_L} V_m = -V_L \quad (26)$$

We define two new POVMs $s(\pm)$ with exactly the same elements and basic weights as the original POVM except for the following changes:

$$p_{m \neq L}^1(\pm) = p_m^1 \pm \delta \frac{\lambda_m}{\lambda_L} \quad ; \quad p_L^2(\pm) = p_L^2 \pm \delta \quad (27)$$

The two new POVMs average to give the original, are complete by construction, and if the δ is chosen small enough, will not violate any of the homogeneous constraints or positivity. The same argument can easily be extended to the case where more than one of the vectors from the set is placed in M_2 . Therefore we require that any set of linearly dependent basic vectors contains non-zero weights in at most one fixed element of an extremal POVM $\square$.

The exact utility of the basic vectors approach will become clearer in the examples that we tackle later, where we will explicitly construct sets of basic vectors. Unfortunately it is not always easy to do this construction. Nevertheless, there is one approach based upon construction of two outcome extremal POVMs that can allow us to draw some interesting general conclusions. Let us consider the set of feasible two outcome POVMs:

$$\{M, \mathbf{1} - M\}. \quad (28)$$

Each such measurement is characterised by just one element M , as the other is fixed by completeness. Suppose then that we can somehow find the extreme points of this set, and they are given by the set $\{M(\alpha)\}$, where α represents some parameters (discrete or continuous) that label the extrema. Then it is clear that $\{M(\alpha)\}$ will form a set of basic vectors for feasible POVMs with *any* number of outcomes. This is because any particular element M_i of an N -outcome POVM can be viewed as a member of a two outcome POVM $M_i, \mathbf{1} - M_i$, where the remaining elements have been grouped together as one outcome, and so any element of a feasible measurement can always be written as a convex combination of the $\{M(\alpha)\}$. This feature will be particularly useful to us in the construction of the separable OO symmetric extrema, as in that case we will be able to construct the two outcome extrema more easily.

There are also more general conclusions that can be made from this observation. Suppose that each two-outcome feasible POVM element can be constrained to live on a K -dimensional vector space. If the elements are $m \times m$ Hermitian operators, then there is a natural upper bound to K of m^2 , although this can be reduced in cases of symmetry. Then by Lemma 7 above, we know that each linearly dependent set of

basic vectors can contribute non-zero weights to at most one POVM element. As any $K + 1$ elements drawn from $\{M(\alpha)\}$ are linearly dependent, we must conclude that in any K non-zero outcome extremal feasible measurement, each element must be proportional to one and only one distinct member of $\{M(\alpha)\}$. Hence we have that:

Lemma 8. In any feasible extremal measurement of K non-zero outcomes, each POVM element must be proportional to an element of a 2-outcome extremal feasible POVM.

This result is interesting as it shows how information from 2-outcome optimisations may be used to draw conclusions about problems with higher numbers of outcomes. Note that the possible constants of proportionality will be uniquely fixed by completeness, as each of the K 2-outcome extrema contributing to the POVM are linearly independent. It is important to note that there cannot be any extremal feasible measurements with more than K non-zero outcomes anyway by the linear independence requirement.

We are now in a position to apply these tools to the cases of symmetry that we will consider in the rest of the paper.

VI. BELL AND OO SYMMETRIC MEASUREMENTS

In the previous section we have discussed a number of useful results that we can use calculate the extreme points of separable POVMs. Here we will now use these techniques and notations to derive the set of local measurements of Bell and OO symmetries, and to characterise the extreme points.

The symmetry groups considered here are all taken from the examples discussed in [22]. They have the convenient property that the invariant POVMs have elements that can be written as a decomposition into a finite number n of orthogonal projectors $A, B, C, \dots$:

$$M_i = a_i A + b_i B + c_i C, \dots \quad (29)$$

(i.e. the *commutant* [35] of the group is abelian). Therefore each POVM element can be represented by a vector of coefficients $(a_i, b_i, c_i, \dots)^T$. As at most n vectors with n components can be linearly independent, this means that we will be able to write the most general extremal PPT POVM for the symmetry group under consideration as an $n \times n$ matrix, where each column j contains the column vector $(a_j, b_j, c_j, \dots)^T$ corresponding to the POVM element M_j .

Another key feature of the symmetries that we consider is that checking the positivity of the partial transposition of a particular POVM element is relatively simple, as the partial transposition of each invariant POVM element can always be written in the following way:

$$M_i^\Gamma = a'_i A' + b'_i B' + c'_i C', \dots \quad (30)$$

where the $(a'_i, b'_i, c'_i, \dots)$ form a vector of real coefficients and the $(A', B', C', \dots)$ are a set of mutually orthogonal projectors.

Note that this is not always the case, in that there can be local symmetry groups with an abelian commutant whose partial transposition is no longer abelian. Nevertheless, there are many interesting symmetries that have this property, including those of Bell diagonal states that we consider next.

A. Local observation with Bell symmetries

In this section we will derive the extreme points of the set of Bell Diagonal PPT POVMs, and then see that they can be attained locally. We are forced to adopt this route, as no direct and simple solution in the manner of the Isotropic/ Werner case is possible for the Bell group. A more detailed explanation for this will be given in the appendix.

First we will explain our notation. The Bell basis is defined as the orthonormal set of two qubit pure states:

$$\begin{aligned} |\Phi^\pm\rangle &= \frac{1}{\sqrt{2}}(|00\rangle \pm |11\rangle) \\ |\Psi^\pm\rangle &= \frac{1}{\sqrt{2}}(|01\rangle \pm |10\rangle), \end{aligned} \quad (31)$$

and it constitutes a set of projectors that is invariant under the symmetry group $\{\sigma_x \otimes \sigma_x, \sigma_y \otimes \sigma_y, \sigma_z \otimes \sigma_z, \mathbf{1} \otimes \mathbf{1}\}$ discussed in [22]. Consequently, any POVM element invariant under the same group can be written as:

$$\begin{aligned} M_i &= a_i |\Psi^+\rangle\langle\Psi^+| + b_i |\Psi^-\rangle\langle\Psi^-| \\ &+ c_i |\Phi^+\rangle\langle\Phi^+| + d_i |\Phi^-\rangle\langle\Phi^-|. \end{aligned} \quad (32)$$

As we require linear independence between the non-zero POVM elements, we can restrict our attention to candidate extremal measurements represented by a 4×4 matrix, where each column $i, i = 1 \dots 4$ has elements given by $\{a_i, b_i, c_i, d_i\}^T$, the quadruple representing the POVM element M_i :

$$s = \begin{pmatrix} a_1 & a_2 & a_3 & a_4 \\ b_1 & b_2 & b_3 & b_4 \\ c_1 & c_2 & c_3 & c_4 \\ d_1 & d_2 & d_3 & d_4 \end{pmatrix}. \quad (33)$$

In each column all elements must be non-negative, and each row must sum to 1 for completeness.

Definition 9. A largest element in a given column will be referred to as the *maximal* element of the column. In many cases the maximal element of a column will not be unique, in which case we are free to choose among the possibilities.

The condition that each column corresponds to a PPT measurement is that a maximal element of a column must be less than or equal to the sum of the other three elements of the same column.

Definition 10. We will refer to a column as being *tight* if it has a maximal element *equal* to the sum of the remaining matrix elements in its column.

Lemma 11. In an extremal Bell POVM with a finite integer N of non-zero outcomes, at least $N - 1$ columns must be tight.

Proof. Suppose that we consider two columns, say without loss of generality the first two. Suppose that there are two rows such that both columns contain non-zero matrix elements. Suppose w.l.o.g. these two rows are the top two rows. Then we can perturb the POVM as follows with a small positivity δ :

$$s(\pm) = s \pm \begin{pmatrix} +\delta & -\delta & 0 & 0 \\ +\delta & -\delta & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (34)$$

If δ is small enough, this perturbation maintains positivity and completeness. Therefore PPT-ness must be violated. Computing the partial transpose of M_i in eq. 32 the positivity of M_i^Γ is equivalent to

$$a_i + b_i \geq |c_i - d_i| \quad (35)$$

$$c_i + d_i \geq |a_i - b_i|. \quad (36)$$

If PPT-ness should be violated by arbitrarily small perturbations δ this implies that at least one of these conditions has to be an equality. Together with the positivity of the a_i, b_i, c_i and d_i , this implies the tightness of at least one column.

Alternatively, it could be the case that there are no two rows in which both columns contain non-zero matrix elements. This can only be the case if one of the columns contains only two non-zero matrix elements. In that case both non-zero matrix elements of the column in question must be maximal in order for it to be PPT, and so that column must be tight.

We can take any two columns in this argument, and so at least $N - 1$ of the non-zero columns must be tight $\square$.

Lemma 12. Any tight column j can itself be written in the following way:

$$\sum_i p_i^j P_i \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + q^j \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad (37)$$

where the summation runs over all permutation matrices P_i and the coefficients $\{p_i^j, q^j\}$ form a probability distribution.

Proof. Let the sum of the matrix elements in the column be $2\sigma_j$. Then as the column is tight, we must have that its maximal element is equal to σ_j . Therefore the column is majorised [36] by the column vector:

$$\begin{pmatrix} \sigma_j \\ \sigma_j \\ 0 \\ 0 \end{pmatrix}. \quad (38)$$

It is a well known result that if a column vector $\vec{g}$ majorises a column vector $\vec{h}$ then $\vec{h}$ can be written as a convex combination of permutations of $\vec{g}$ [36]. Hence we can write that:

$$\sigma_j \sum_i x_i^j P_i \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + (1 - \sigma_j) \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \quad (39)$$

where the $\{x_i^j\}$ form a probability distribution. As no maximal element can be greater than 1 (from completeness), the $\sigma_j, 1 - \sigma_j$ also form a probability distribution, and so by setting $p_i^j = \sigma_j x_i^j$ and $q^j = \sigma_j$ we have a decomposition of the column as stated in the lemma. Note that the decomposition need not be unique.

Lemma 13. All columns j in a Bell PPT POVM can be expressed as

$$\sum_i p_i^j P_i \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + q^j \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} + r^j \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad (40)$$

where the $\{p_i^j, q^j, r^j\}_i$ form a probability distribution.

Proof. Let us for now just consider extremal measurements of two non-zero outcomes. One of the two non-zero columns of any such measurement must be tight, let this w.l.o.g. be column 1. Then from completeness and Lemma 12 we can expand column 2 as:

$$\begin{aligned} \begin{pmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{pmatrix} &= \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} - \left[\sum_i p_i^1 P_i \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + q^1 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \right] \\ &= \sum_i p_i^1 P_i \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix} + q^1 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \\ &:= \sum_i p_i^2 P_i \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix} + r^2 \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \end{aligned}$$

where as before $\{p_i^2, r^2\}$ also forms a probability distribution. Hence the theorem holds for all 2 non-zero outcome extremal PPT Bell measurements, and by convexity holds for the non-extremal ones as well. But then any column from an N outcome PPT measurement can be viewed as a member of a two outcome measurement where the remaining columns have been summed. Therefore the result is true for any column in any Bell PPT POVM. $\square$.

A consequence of this lemma is that the set of column vectors of the form:

$$P_i \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} \text{ or } \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix} \quad (41)$$

forms a set of basic vectors for the Bell Diagonal PPT POVMs. For a column j , the probability p_i^j or r^j will hence

be taken as the weights of the corresponding basic vector. We will refer to as $(1, 1, 1, 1)^T$ as the *identity basic vector* as it corresponds to the identity POVM.

Lemma 14. There are no 4 non-zero outcome extremal Bell POVMs.

Proof. In a 4 non-zero outcome extremal POVM, each non-zero column must have weight from only one basic vector distinct from the basic vectors contributing to the other columns. Otherwise the linear independence condition in Lemma 7 will be violated (as any five 4-component column vectors must be linearly dependent).

If one column contains weight from the identity, then the POVM cannot be extremal, as it can be expressed as the convex sum of the identity and a renormalised POVM that consists of only the remainder from the original.

For four outcomes where no column contains weight from the identity, a little thought shows that the only possibility that can maintain completeness is a POVM given by a row and/or column permutations of the following:

$$\begin{pmatrix} w & 0 & 0 & z \\ w & x & 0 & 0 \\ 0 & x & y & 0 \\ 0 & 0 & y & z \end{pmatrix}, \quad (42)$$

However, this measurement cannot be extremal as the columns are linearly dependent. To see this, note that we can obtain the zero vector by (a) multiplying each column by a non-zero factor such that each non-zero matrix element has the same value, and then (b) subtracting the 2nd and 4th columns from the sum of the 1st and 3rd. Therefore, there are no four non-zero outcome extremal PPT Bell POVMs.

Lemma 15. There are no 3 non-zero outcome extremal Bell POVMs.

Proof. As with the proof for four non-zero outcomes, there can be no column with weight from the identity. Therefore we require that each column can be decomposed into convex sums of basic vectors other than the identity, such that no more than four basic vectors are involved in total. Therefore two columns must be directly proportional to a basic vector, as each basic vector can have non-zero weight in at most one column. Let these two columns be 1 and 2 w.l.o.g.. There are essentially only two possibilities upto row and column permutations - the first two columns have a common row in which they both have non-zero matrix elements, or they have no such common row. The third column is fixed by completeness. Therefore up to row and column permutations we have:

$$\begin{pmatrix} x & 0 & 1-x & 0 \\ x & y & 1-x-y & 0 \\ 0 & y & 1-y & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \text{ or } \begin{pmatrix} x & 0 & 1-x & 0 \\ x & 0 & 1-x & 0 \\ 0 & y & 1-y & 0 \\ 0 & y & 1-y & 0 \end{pmatrix}, \quad (43)$$

but these POVMs cannot be extremal unless they have one further zero column. For the first matrix we can see this from the following. Consider the perturbation

$$\pm \delta \begin{pmatrix} +1 & 0 & -1 & 0 \\ +1 & -1 & 0 & 0 \\ 0 & -1 & +1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (44)$$

Unless either x or y are zero, this perturbation maintains completeness, positivity and PPT-ness. Hence the first matrix cannot be extremal. For the second matrix we can consider the following perturbation:

$$\pm \delta \begin{pmatrix} +1 & 0 & -1 & 0 \\ +1 & 0 & -1 & 0 \\ 0 & -1 & +1 & 0 \\ 0 & -1 & +1 & 0 \end{pmatrix}. \quad (45)$$

This perturbation implies that to be extremal either x or y are zero, or that the third column is tight. We cannot have three non-zero columns if x or y are zero, so let us deal with the possibility that the third column is tight. The third column can only be tight if one of x or y is 1. Suppose w.l.o.g. that $x = 0$, then the second and third columns become proportional, and so it cannot be extremal. Hence neither of the possibilities in (43) can be extremal $\square$.

Therefore there the only extremal POVMs of Bell diagonal form are those of at most two non-zero outcomes.

Theorem 16. The extremal points of the PPT Bell diagonal POVMs can be obtained by local protocols. The extremal points are given by the possible row and column permutations of the following POVMs:

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}; \quad \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}. \quad (46)$$

Proof. The fact that these are the only extremal points can be proven using similar arguments to the ones above. They are also locally attainable, as the first matrix corresponds to the ‘do nothing’ measurement, and the other POVMs correspond to POVMs of the form:

$$M = \sigma_k \otimes \sigma_l [|00\rangle\langle 00| + |11\rangle\langle 11|] \sigma_k^\dagger \otimes \sigma_l^\dagger; \quad \mathbb{1} - M \quad (47)$$

where the σ_i are drawn from the four Pauli matrices. These POVMs are trivially locally attainable. They simply correspond to measurements in a product basis, and can be achieved using one way communication $\square$.

Despite some prior results showing that the four orthogonal Bell states cannot be perfectly discriminated locally [19,16], this result is interesting as it shows that there can be a huge difference between locally and globally obtainable information under *all* figures of merit, even when we are only dealing with two qubits. Indeed, as the extremal local measurements involve only 2 non-zero outcomes, there can be at most 1 classical bit of information obtained by local measurements of Bell diagonal states. However, in the global case there can

be 4 orthogonal Bell diagonal states that can be perfectly discriminated, thereby allowing a full 2 classical bits of information to be obtained globally. It might be tempting to propose a bit hiding scheme on the basis of this observation. However, the scheme would be susceptible to cheating with a prior shared singlet state (as then Alice could teleport her particle to Bob), and hence the technology required to break the scheme is equivalent to that required to implement it.

B. Local Observations with OO Symmetries

In the notation of [22], the OO group is the set of unitaries of the form $O \otimes O$, where O is an orthogonal transformation (i.e. $OO^T = I_{A/B}$) in some fixed local bases. Any POVM element invariant under this group can be written in the form:

$$a[|+\rangle\langle+|] + b\left[\frac{(\mathbb{1} - F)}{2}\right] + c\left[\frac{(\mathbb{1} + F)}{2} - |+\rangle\langle+|\right],$$

where $F = \sum |ij\rangle\langle ji|$ is the swap operator, a, b, c are real coefficients, and the three terms in square brackets are mutually orthogonal projections. Hence a full measurement obeying the OO symmetry with N outcomes is characterised completely by the triples (a_k, b_k, c_k) , corresponding to each of the POVM elements indexed by $k = 1, \dots, N$. One could try to look for local protocols to match these triples in a similar spirit to the solution presented for the Isotropic case. To be specific, it might be hoped that a linear invertible transformation could be found taking the (a_k, b_k, c_k) to new triples (x_k, y_k, z_k) , such that the (x_k, y_k, z_k) are the positive coefficients of some local protocol iff the original (a_k, b_k, c_k) correspond to a PPT POVM. However, we can show that no such naive transformation exists for the OO symmetries (see appendix 1 for details). Consequently, in the following we will instead restrict our attention to the *extreme points* of the convex set of OO symmetric PPT POVMs. We will present local protocols that attain these extreme points, and thereby argue that all OO symmetric PPT POVMs can be attained locally. First we need to show how to construct these extreme points.

1. Extreme points of the OO symmetric POVMs

First let us discuss the partial transposition of OO symmetric operators. Partial transposition takes M to an operator of the same form, with new coefficients given by:

$$\begin{pmatrix} a' \\ b' \\ c' \end{pmatrix} = R \begin{pmatrix} a \\ b \\ c \end{pmatrix}. \quad (48)$$

where

$$R = \frac{1}{2d} \begin{pmatrix} 2 & d(1-d) & (d+2)(d-1) \\ -2 & d & (d+2) \\ 2 & d & (d-2) \end{pmatrix}. \quad (49)$$

In order to satisfy the constraints of being a PPT POVM, all the triples (a_k, b_k, c_k) corresponding to each element M_k must satisfy:

$$\begin{aligned} a_k, b_k, c_k &\geq 0 \text{ (positivity)} \\ a'_k, b'_k, c'_k &\geq 0 \text{ (PPT-ness)} \\ \sum_k a_k &= \sum_k b_k = \sum_k c_k = 1 \text{ (normalisation)}. \end{aligned} \quad (50)$$

We would like to construct a set of basic vectors for the OO symmetries. One way to do this is to first construct the set of 2-outcome PPT extrema under these symmetries. In such situations given one element M the other is fixed as $\mathbb{1} - M$ by completeness. We hence require:

$$\begin{aligned} \mathbb{1} &\geq M \geq 0 \\ \mathbb{1} &\geq M^T \geq 0. \end{aligned}$$

Denoting the only free element by a column vector of its triple $M = (a, b, c)^T$, the constraints will result in a polyhedron in R^3 , for which the extreme points are the vertices (see figure 1).




FIG. 1. The polyhedron of allowed PPT 2-outcome POVMs under OO symmetry, sketched for $d = 3$. The vertices are given in equation (51).

It is trivial but tedious to calculate these vertices. They are given by the following POVM elements:

$$\begin{aligned} A1 &: \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} & A2 &: \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \\ B1 &: \begin{pmatrix} 0 \\ 0 \\ 2d/[d+2][d-1] \end{pmatrix} \\ B2 &: \begin{pmatrix} 1 \\ 1 \\ (d+1)(d-2)/[d+2][d-1] \end{pmatrix} \end{aligned}$$

$$\begin{aligned}
C1 : & \begin{pmatrix} 1 \\ 1/[d-1] \\ (d-2)/[d+2][d-1] \end{pmatrix} \\
C2 : & \begin{pmatrix} 0 \\ (d-2)/[d-1] \\ d^2/[d+2][d-1] \end{pmatrix} \\
D1 : & \begin{pmatrix} 1 \\ 0 \\ 2/[d+2] \end{pmatrix} \quad D2 : \begin{pmatrix} 0 \\ 1 \\ d/[d+2] \end{pmatrix}
\end{aligned} \tag{51}$$

Those POVM elements labelled with the same upper case letter are *complementary*, in the sense that they sum to the identity. Hence each letter labels a complete extremal PPT *OO* symmetric POVM of 2 non-zero outcomes. The entire set of vectors in equation (51) hence also forms a set of basic vectors for *OO* symmetries. We would like to use this set to construct the entire set of extremal points. We require any candidate extremal POVM to consist of measurement elements that are linearly independent. Therefore we need only restrict our attention to POVMs with at most 3 non-zero outcomes, i.e. those measurements that can be represented by a 3×3 matrix:

$$s = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}, \tag{52}$$

where each column characterises one POVM element. As a consequence of Lemma 7, we can only use three of our basic vectors to construct the POVM, and as we require all three columns to be non-zero, each column of the extremal 3-outcome POVMs must be *proportional* to one and only one distinct basic vector. As there are essentially only 7 important basic vectors to choose from in the set above ($A1$ is trivial), we are already constrained as to the possible form. However, there is another observation that will provide much tighter constraints:

Lemma 17. It is not possible to have 2 basic vectors from the same complementary pair contributing non-zero weights in the same extremal POVM. Consequently, no extremal POVM can have any column with non-zero weight from the identity basic vector, unless it is the trivial POVM with only one non-zero outcome.

Proof. Let us suppose that the two complementary basic vectors are V_1 and V_2 respectively. Then suppose that two POVM elements j, k (where we can have $j = k$) contain weights p_1^j and p_2^k . Suppose that $p_1^j \leq p_2^k$. Then we can write the whole POVM as a convex combination with probability p_1^j of a two-outcome POVM ($M_1 = V_1, M_2 = V_2$) and $1-p_1^j$ of the rest of the POVM rescaled to be complete. Hence the POVM cannot be extremal unless one of the p_1^j, p_2^k is zero $\square$.

An immediate consequence of Lemma 17 is that we cannot use either of the basic vectors $A1, A2$ to construct our POVM. Another consequence is that from the remaining possible basic vectors, we must utilise exactly one vector from each of the pairs $\{B1, B2\}$, $\{C1, C2\}$ and $\{D1, D2\}$. We also require that in choosing each basic vector, and its weight in a

column, we must respect completeness. Therefore it is not possible to choose all three selected basic vectors with a 1 in the top component. This leaves only six possible choices for the basic vectors with non-zero weight:

$$\begin{aligned}
& \{B1, C1, D1\} \\
& \{B1, C1, D2\} \\
& \{B1, C2, D1\} \\
& \{B2, C1, D2\} \\
& \{B2, C2, D1\} \\
& \{B2, C2, D2\}
\end{aligned} \tag{53}$$

It is not difficult, although tedious, to verify that the only combination that can be made complete with positive weights is the combination $\{B1, C1, D2\}$. We recommend the use of a standard computational package for algebraic manipulation, with which it can readily be verified that the selection $\{B2, C1, D2\}$ is linearly dependent, and that the remaining possibilities other than $\{B1, C1, D2\}$ all require negative weights in order to satisfy completeness [37]. For $\{B1, C1, D2\}$ the weights are also fixed uniquely, as the solution involves a set of 3 linearly independent equations in 3 unknowns. It turns out that two of the three weights are 1 (those of $B1$ and $C1$), and as these elements are also 2-outcome extrema, they cannot be perturbed in any way while keeping the whole POVM feasible. Hence this whole 3-outcome POVM is extremal, and we have the following:

Lemma 18. The only genuine 3-outcome extremal POVM for *OO* symmetries, upto relabelling the outcomes, is given by the POVM with elements represented by the following triples:

$$\begin{aligned}
M_1 &= (0, 0, \frac{2d}{(d+2)(d-1)}) \\
M_2 &= (0, \frac{d-2}{d-1}, \frac{d(d-2)}{(d+2)(d-1)}) \\
M_3 &= (1, \frac{1}{d-1}, \frac{d-2}{(d+2)(d-1)}),
\end{aligned} \tag{54}$$

where we have made the slight abuse of notation that a POVM element will be set equal to the triple representing it.

2. Locally attaining the extreme points of the *OO* symmetric POVMs

In trying to attain the extreme points derived above, we will first consider only the 2-outcome extrema, given in equation (51). The important question is how to realise the extreme points of this polyhedron using LOCC measurements. The measurement A and its complement are trivial, they correspond to the do nothing POVM. The measurement D and its complement are also not too difficult to obtain. It is easy to confirm that if Alice and Bob carry out an ‘orthogonal twirl’ [31,38] followed by the locally implementable projection:

$$\sum_{i=0}^{d-1} |ii\rangle\langle ii| \quad (55)$$

then they will achieve the POVM D , and so both D and its complement can be attained.

The points B , C , and complements are a little more subtle. However, by performing an orthogonal twirl [38] it is not difficult to show that if we can find a set of overcomplete pure states on Alice's particle $\{|q\rangle\}$, $q = 1 \dots L$, $L \geq d$ with the properties that:

$$\frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| = I_A \quad (56)$$

$$\forall q \quad \text{tr}\{|q\rangle\langle q|(|q\rangle\langle q|)^T\} = 0 \quad (57)$$

then the following local measurements:

$$\frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| \otimes |q\rangle\langle q| \quad (58)$$

and

$$\frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| \otimes (|q\rangle\langle q|)^T \quad (59)$$

will attain points B and C respectively when preceded by the orthogonal twirl. It remains to show that a set of pure states satisfying equations (56) and (57) exists. Two pure states with property (57) are:

$$\begin{aligned} |u\rangle &:= 1/\sqrt{2}(|r\rangle + i|s\rangle) \\ |v\rangle &:= 1/\sqrt{2}(|r\rangle - i|s\rangle) \end{aligned} \quad (60)$$

for any two computational basis states $|r\rangle, |s\rangle$. Suppose that d is even. Then Alice can obtain a complete set of $\{|q\rangle\}$ satisfying the properties by partitioning her space in 2-level orthogonal subspaces, and constructing a $|u\rangle, |v\rangle$ for each subspace. If d is odd, then she can do the same until she is left with a 3-level subspace. We therefore need to construct such a set of pure states for this 3-level subspace. One route to a solution is as follows. Without loss of generality assume that this residual subspace is spanned by computational basis states $|1\rangle, |2\rangle, |3\rangle$. Then Alice can pick any irreducible representation of a finite group in 3×3 real orthogonal matrices, i.e. $OO^T = I$. Suppose this set is $\{O_m | m = 1 \dots L\}$. Then the set of pure states:

$$O_m[1/\sqrt{2}(|1\rangle + i|2\rangle)] \quad (61)$$

will satisfy the requirements (56) and (57). It seems likely that this solution for odd d is overly complicated, however, we were not able to find any significant simplifications.

Therefore we see that all extreme points of the set of 2-outcome PPT OO symmetric measurements can be obtained by local protocols involving only one way communication. Now we need to turn to the 3-outcome element presented

in equation (54). Given the existence of a set of pure states $\{|q\rangle | q = 1 \dots L, L \geq d\}$ satisfying the conditions (56) and (57), it is not too difficult to verify that the local POVM defined by the following elements:

$$\begin{aligned} N_1 &= \frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| \otimes |q\rangle\langle q| \\ N_2 &= \frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| \otimes (|q\rangle\langle q|)^T \\ N_3 &= \frac{d}{L} \sum_{q=1}^L |q\rangle\langle q| \otimes [\mathbf{1}_B - |q\rangle\langle q| - (|q\rangle\langle q|)^T] \end{aligned} \quad (62)$$

achieves the M_1, M_2, M_3 of eq.(54) respectively when preceded by the orthogonal twirl.

Therefore all extreme points for N-outcome OO symmetric POVMs can be attained by local protocols.

VII. SUMMARY & DISCUSSION

The classification of Separable and PPT operations seems to be particularly suited to the question of measurements, where we find that a POVM can be implemented by a separable operation iff its elements are separable, and by a PPT operation iff its elements are PPT. As the Separable/PPT POVMs are mathematically more tractable than the LOCC measurements, and as we have the (strict) inclusions $\text{LOCC} \subset \text{Separable} \subset \text{PPT} \subset \text{Global}$, these classes of operation are useful for deriving bounds on what we can achieve with LOCC observations. For this reason it is useful to have techniques for deriving extremal measurements under the constraints of separability and PPT-ness. In this context we have discussed a number of useful tools. The partial transposition isomorphism is particularly useful for PPT measurements under partially conjugated symmetries, whereas the linear independence requirements and basic vectors approach are more generally useful for extremising POVMs under any homogeneous constraints. Applying these techniques we were able to show that for Isotropic, Werner, Bell and OO symmetries, the separability of POVM elements is necessary and sufficient for local implementability. There is no obvious reason why this is the case, as it is clear from the results of [30] that not all separable POVMs can be implemented locally.

Perhaps the most immediate consequences of our results are for local state discrimination. Imagine that we have been given one of i states ρ_i with probability p_i . What is the best way to tell which state we have been given? Questions such as this are significant for entanglement distillation protocols, where 'target' states are often locally measured to give information about 'source' states [10,13], and also for cryptographic schemes. In such scenarios one expects the extremal local POVMs to be the optimal possible measurements for most reasonable cost functions, regardless of the number of

states or prior probabilities. Therefore our results also essentially give optimal protocols for the local discrimination of Isotropic, Werner, Bell and *OO* states. In the context of state discrimination the Bell symmetries give the largest (in general) separation between local and global state discrimination. If we have global access to one of the four orthogonal Bell pure states, then we can discriminate them perfectly, potentially obtaining 2 classical bits of information. However, if we are restricted to acting locally, the best measurement we can do has at most 2 non-zero outcomes, showing that we can locally obtain at most 1 bit of information. Unfortunately this does not supply an interesting bit hiding scheme in the manner of [9,21], as the technology required to implement the scheme is the same as the technology required to break it - Alice and Bob can share prior Bell states and use them to teleport Alice's state to Bob, thereby allowing Bob full access to the state without having to meet Alice. Nevertheless, our results give the first examples of mixed states for which a full quantitative analysis of optimal state discrimination can be performed.

We hope to be able to apply the techniques from this paper to more ambitious symmetries, perhaps considering more specific cost functions or prior probabilities in order to make the calculations more tractable. A fuller understanding of local measurements could have applications not only for local state discrimination and entanglement distillation, but also for any situation in which local observations are required to demonstrate a phenomenon, such as in demonstrations of non-locality, and also the recent program of research on the experimental implementation of entanglement witnesses [14].

VIII. ACKNOWLEDGEMENTS

SV thanks Oliver Rudolph, Adam Brazier and Tilo Eggeling for sharing their insights, and Paolo Lo Presti & Mauro D'Ariano for discussions about global extremal POVMs. We acknowledge financial support from EC project EQUIP (IST-1999-11053), EC project ATESIT (contract IST-2000-29681), EC project ATESIT (contract IST-2000-29681), INFM PRA 2001 CLON, US Army Grant DAAD 19-02-0161, the UK EPSRC and the European Science Foundation Programme on Quantum Information Theory and Quantum Computing.

IX. APPENDIX 1: NO 'NAIVE' SOLUTION POSSIBLE FOR *OO* SYMMETRIES.

In this appendix we will see that no 'naive' solution can be found for *OO* symmetric POVMs with the same simplicity as the solution for the Isotropic/Werner measurements. First we need to clarify what we mean by 'naive'. We will consider local measurement protocols consisting of POVM elements N_k , each of which can be written in the following form:

$$N_k = x_k X + y_k Y + z_k Z \quad (63)$$

where the $\{X, Y, Z\}$ are orthogonal projectors that sum to the identity, and the (x_k, y_k, z_k) are triples of real coefficients. Then the requirements of completeness and positivity mean that the sets of $\{x_k\}$, $\{y_k\}$ and $\{z_k\}$ must form probability distributions. We will also assert that *any such valid POVM is also local*. Under these assertions we will also impose the following requirement:

- Any PPT POVM of *OO* symmetry can be attained by performing an orthogonal twirl followed by a (local) measurement of the stipulated form (63).

We will show that no such solution is possible for the *OO* case, even though a solution with essentially the same features was possible for the Isotropic case.

Let us represent each POVM element N_k by the column vector $\vec{v}_k = (x_k, y_k, z_k)^T$. Then under twirling, N_k will be taken to an *OO* symmetric POVM element. Let this element be represented by the column vector $\vec{w}_k = (a_k, b_k, c_k)^T$, using the same notation as the main text. There will be a linear transformation L , such that $L \cdot \vec{v}_k = \vec{w}_k$ represents the effect of the twirling.

As stated in the main text, the the only non-zero elements in extremal (PPT) POVMs must be linearly independent (Theorem 5). Let us represent the set of general extremal POVMs of the stipulated form (63) by the set $\{r_i\}$, and the extrema of the PPT *OO* symmetric form by the set $\{s_i\}$. As a consequence of the linear independence requirement, every one of these extrema can be expressed as 3×3 matrices, where the columns are linearly independent and represent each of three non-zero POVM elements. Hence two general members $r \in \{r_i\}$ and $s \in \{s_i\}$ of these sets can be written:

$$r = \begin{pmatrix} x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \\ z_1 & z_2 & z_3 \end{pmatrix}, \quad s = \begin{pmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{pmatrix}. \quad (64)$$

As the columns of these matrices are linearly independent, the matrices are invertible. This will be important in the subsequent discussion.

Now let us see the implications of the above assertions for the form of the transformation L . We are forced to have the following properties:

- L must have non-negative matrix elements, as its elements are calculated from the traces of products of non-negative operators.
- Each row of L must sum to 1. This is from the fact that twirling retains completeness of the POVM.
- L must be invertible. As we wish *all* PPT *OO* POVMs to be obtainable from the stipulated protocol, the extreme points $\{s_i\}$ of the PPT *OO* symmetric POVMs must be contained within set obtained from L acting on the extrema $\{r_i\}$ of the stipulated measurement. Consequently for some r, s we have $L \cdot r = s$, and hence $L^{-1} = r \cdot s^{-1}$.

Let us now consider the transformation R of equation (48). The vector:

$$R \cdot L \cdot \vec{v}_k = R \cdot \vec{w}_k \quad (65)$$

gives the partial transpose of the twirled POVM element corresponding to N_k . As we wish *any* PPT POVM be attainable by a local protocol of our postulated form, and as we require that L be invertible, we also require that both:

$$L \cdot \vec{v}_k \geq 0 \quad \forall k \quad (66)$$

and

$$R \cdot L \cdot \vec{v}_k \geq 0 \quad \forall k \quad (67)$$

hold if and only if the sets $\{x_k\}$, $\{y_k\}$ and $\{z_k\}$ form probability distributions. We also must require the following properties for the matrix product $R \cdot L$:

- $R \cdot L$ must contain only non-negative elements. This can be seen as follows. If we pick a three element local measurement characterised by:

$$\vec{v}_1 = (1, 0, 0)^T ; \vec{v}_2 = (0, 1, 0)^T ; \vec{v}_3 = (0, 0, 1)^T \quad (68)$$

then this must produce vectors $\vec{w}_k$ with non-negative partial transposition, and it can readily be verified that this implies that the elements of $R \cdot L$ must be non-negative.

- The rows of $R \cdot L$ must sum to 1. This can be shown from the fact that the same property holds for both R and L .

Now consider the following disallowed choice for the vectors $\vec{v}_k$, where ϵ is small and positive:

$$\vec{v}_1 = (1/3, 1/3, -\epsilon)^T ; \vec{v}_2 = (1/3, 1/3, (1+\epsilon)/2)^T ; \vec{v}_3 = (1/3, 1/3, (1+\epsilon)/2)^T. \quad (69)$$

This choice of vectors does not give a valid choice of measurement vectors, as the first one contains a negative component. However, the vectors $\vec{v}_2$ and $\vec{v}_3$ can in principle come from valid probability distributions by themselves, and so it must be the case that either $L \cdot \vec{v}_1$ or $R \cdot L \cdot \vec{v}_1$ contains a negative component. It can be shown that this, together with the fact that both L and $R \cdot L$ must be row stochastic, implies that at least one row of either L or $R \cdot L$ must be $(0, 0, 1)$. Similarly we can ‘place the ϵ ’ in different rows in the above vectors to show that in fact we require that at least one row of either L or $R \cdot L$ must be $(0, 1, 0)$ and at least one row should be $(1, 0, 0)$. Regardless of where these rows appear, as $R = R^{-1}$ contains some negative matrix elements, no such solution is possible, as both L and $R \cdot L$ are required to have non-negative matrix elements. Although the proof that we have given is tailored to OO symmetries, it can be modified relatively easily for other situations, including bases of Bell diagonal states.

- [1] M.B. Plenio and V. Vedral, *Cont. Phys.* **39**, 431 (1998); R.F. Werner, *Quantum information – an introduction to basic theoretical concepts and experiments*, Springer Tracts in Modern Physics (Springer, Heidelberg, 2001).
- [2] K. Kraus, ‘Lecture Notes: States, Effects and Operations’, Springer.
- [3] J. Eisert, K. Jacobs, P. Papadopoulos and M.B. Plenio, *Phys. Rev. A* **62**, 52317 (2000).
- [4] D. Collins, N. Linden and S. Popescu, *Phys. Rev. A* **64**, 032302 (2001).
- [5] A. Jamiolkowski, *Rep. Math. Phys.* No. 4, **3** 275 (1972).
- [6] W. Dür and J.I. Cirac, *Quant. Inf. Comp.*, Vol.2, No. 3, 240-254 (2002); J.I. Cirac, W. Dür, B. Kraus and M. Lewenstein, *Phys. Rev. Lett.* **86**, 544 (2001).
- [7] E.M. Rains, *Phys. Rev. A* **60**, 173 (1999); **60**, 179 (1999).
- [8] E.M. Rains, *IEEE Trans. Inform. Theory* **47**, 2921, (2001).
- [9] B.M. Terhal, D.P. DiVincenzo and D.W. Leung, *Phys. Rev. Lett.* **86** 5807-5810 (2001).
- [10] C.H. Bennett, D.P. DiVincenzo, J.A. Smolin and W.K. Wootters, *Phys. Rev. A* **54** 3824-3851 (1996); C.H. Bennett, H.J. Bernstein, S. Popescu and W.K. Wootters, *Phys. Rev. A* **53** 2046 (1996).
- [11] V. Vedral, M.B. Plenio, M.A. Rippin, and P.L. Knight, *Phys. Rev. Lett.* **78**, 2275 (1997); V. Vedral and M.B. Plenio, *Phys. Rev. A* **57**, 1619 (1998); M.B. Plenio, S. Virmani and P. Papadopoulos, *J. Phys. A* **33**, L193 (2000); W.K. Wootters, *Phys. Rev. Lett.* **80** 2245-2248 (1998).
- [12] K. Audenaert, M.B. Plenio and J. Eisert, *Phys. Rev. Lett.* **90**, 027901 (2003).
- [13] K.G.H. Vollbrecht and M.M. Wolf, *Phys. Rev. A* **67**, 012303 (2003).
- [14] O. Gühne, P. Hyllus, D. Bruss, A. Ekert, M. Lewenstein, C. Macchiavello and A. Sanpera, *quant-ph/0205089* and *quant-ph/0210134*; P. Horodecki and A. Ekert, *Phys. Rev. Lett.* **89**, 127902 (2002); G. M. D’Ariano, C. Macchiavello and M. G. A. Paris, *quant-ph/0211146*; J.M.G. Sancho and S.F. Huelga, *Phys. Rev. A* **61**, 042303 (2000).
- [15] J. Walgate, A.J. Short, L. Hardy and V. Vedral, *Phys. Rev. Lett.* **85**, 4972 (2000).
- [16] J. Walgate and L. Hardy, *Phys. Rev. Lett.* **89**, 147901 (2002); A. Sen De and U. Sen *quant-ph/0203007*.
- [17] S. Virmani, M. Sacchi, M.B. Plenio and D. Markham, *Phys. Lett. A* **288**, 62 (2001); Yi-Xin Chen and D. Yang, *Phys. Rev. A* **65**, 022320 (2002); Yi-Xin Chen and D. Yang, *Phys. Rev. A* **64**, 064303 (2001); S. Ghosh, G. Kar, A. Roy, D. Sarkar, A. Sen De and U. Sen, *Phys. Rev. A* **65**, 062307 (2002); M. Hillery and J. Mimih, *quant-ph/0210179*.
- [18] M. Horodecki, A. Sen De, U. Sen and K. Horodecki, *Phys. Rev. Lett.* **90**, 047902 (2003).
- [19] S. Ghosh, G. Kar, A. Roy, A. Sen De and U. Sen, *Phys. Rev. Lett.* **87**, 277902 (2001).
- [20] D.P. DiVincenzo, D.W. Leung and B.M. Terhal, *IEEE Trans. Inform. Theory* **48**, 580 (2002); D.P. DiVincenzo, P. Hayden and B.M. Terhal, *quant-ph/0210053*.

- [21] T. Eggeling and R.F. Werner, Phys. Rev. Lett. **89**, 097905 (2002).
- [22] K.G.H. Vollbrecht and R.F. Werner, Phys. Rev. A. **64** 062307 (2001).
- [23] K. Audenaert, J. Eisert, E. Jané, M.B. Plenio, S. Virmani and B. DeMoor, Phys. Rev. Lett. **87**, 217902 (2001); K. Audenaert, B. De Moor, K.G.H. Vollbrecht and R.F. Werner, Phys. Rev. A **66**, 032310 (2002).
- [24] B. Groisman and B. Reznik, Phys. Rev. A **66**, 022110 (2002).
- [25] D. Beckman, D. Gottesman, M.A. Nielsen and J. Preskill, Phys. Rev. A **64**, 052309 (2001).
- [26] R.F. Werner, Phys. Rev. A **40**, 4277 (1989).
- [27] A. Peres, Phys. Rev. Lett. **77** 1413-1415 (1996).
- [28] In [20] this observation was credited to unpublished work by Eric Rains. Hence here we supply a brief indication of a possible proof. Given a PPT POVM $\{M_i\}$ define the CP map:

$$\rho \rightarrow \sum_i \text{tr}(M_i \rho) |i\rangle\langle i| \otimes |i\rangle\langle i| \quad (70)$$

where $\{|i\rangle\}$ is a basis of pure orthogonal states. This is a PPT operation, and leads to the implementation of $\{M_i\}$ if it is followed by a local von Neumann measurement by Alice and Bob in the basis $\{|i\rangle\}$.

- [29] C.H. Bennett, D.P. DiVincenzo, T. Mor, P.W. Shor, J.A. Smolin and B.M. Terhal, Phys. Rev. Lett. **82** (1999) 5385.
- [30] C.H. Bennett, D.P. DiVincenzo, C.A. Fuchs, T. Mor, E. Rains, P.W. Shor and J.A. Smolin, Phys. Rev. A **59**, 1070 (1999).
- [31] The *Isotropic twirling* operation T in $d \times d$ dimensions, $d > 2$, is defined as $T(Q) = \int d\mu_U (U \otimes U^*) Q (U \otimes U^*)^\dagger$, where the integral is performed with respect to the Haar measure,

$U \in U(d)$. The analogous operations for the Bell, Werner and *OO* symmetries will be referred to as the *Bell twirl*, *Werner twirl* and *Orthogonal twirl* respectively.

- [32] See entry for the *Krein-Milman* theorem, in *Collins Dictionary of Mathematics*, Collins (1989). For more detail see Section 18, *Convex Analysis*, R. T. Rockafellar, Princeton (1970).
- [33] G. M. D'Ariano and P. Lo Presti, quant-ph/0301110.
- [34] There is at least one other computable separability criterion known to be independent of PPT that also gives homogeneous constraints, namely the criterion first proposed by Rudolph (O. Rudolph, quant-ph/0202121). Hence the techniques discussed may yield further bounds on LOCC observations using that criterion instead of PPT.
- [35] The commutant of a group g of unitaries is defined as the set of all operators A for which $[U, A] = 0$ for all $U \in g$.
- [36] Theorem **4.3.33**, *Matrix Analysis*, R.A. Horn and C.A. Johnson, Cambridge University Press (1985).
- [37] A *Maple 7* script for this task is available by emailing a request to s.virmani@ic.ac.uk.
- [38] Under orthogonal twirling the product operator $A \otimes B$ becomes an *OO* symmetric operator with the following values for the triple (a, b, c) :

$$\begin{aligned} a &= \frac{\text{tr}\{AB^T\}}{d} \\ b &= \frac{1}{d(d-1)} [\text{tr}\{A\}\text{tr}\{B\} - \text{tr}\{AB\}] \\ c &= \frac{2}{(d+2)(d-1)} \left[\frac{\text{tr}\{A\}\text{tr}\{B\} + \text{tr}\{AB\}}{2} - \frac{\text{tr}\{AB^T\}}{d} \right] \end{aligned}$$